



MarLIN

Marine Information Network

Information on the species and habitats around the coasts and sea of the British Isles

A tubeworm (*Spirobranchus triqueter*)

MarLIN – Marine Life Information Network
Biology and Sensitivity Key Information Review

Karen Riley & Susie Ballerstedt

2005-10-26

A report from:

The Marine Life Information Network, Marine Biological Association of the United Kingdom.

Please note. This MarESA report is a dated version of the online review. Please refer to the website for the most up-to-date version [<https://www.marlin.ac.uk/species/detail/1794>]. All terms and the MarESA methodology are outlined on the website (<https://www.marlin.ac.uk>)

This review can be cited as:

Riley, K. & Ballerstedt, S. 2005. *Spirobranchus triqueter* A tubeworm. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom.
DOI <https://dx.doi.org/10.17031/marlin.sp.1794.2>



The information (TEXT ONLY) provided by the Marine Life Information Network (MarLIN) is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 2.0 UK: England & Wales License. Note that images and other media featured on this page are each governed by their own terms and conditions and they may or may not be available for reuse. Permissions beyond the scope of this license are available [here](https://www.marlin.ac.uk). Based on a work at www.marlin.ac.uk

(page left blank)



Calcareous tubes of two keel worms among several barnacles.

Photographer: David Fenwick

Copyright: David Fenwick Snr.

See online review for
distribution map

Distribution data supplied by the Ocean
Biogeographic Information System (OBIS). To
interrogate UK data visit the NBN Atlas.

Researched by	Karen Riley & Susie Ballerstedt	Refereed by	This information is not refereed.
Authority	(Linnaeus, 1758)		
Other common names	-	Synonyms	<i>Pomatoceros triqueter</i>

Summary

Description

The calcareous tube of *Spirobranchus triqueter* is 3.5 mm wide and up to 25 mm long. It is white, smooth and irregularly curved with a single, median ridge that ends in a projection over the anterior opening. The operculum bears a shallow, dish-shaped plug (ampulla) which is often conical distally, and may have projections on the crown. The colouration of the body is bright but variable, and the crown of tentacles (radioles) are banded with various colours.

Recorded distribution in Britain and Ireland

Common and widespread on all coasts.

Global distribution

Occurs from the coasts of north west Europe to the Mediterranean.

Habitat

Spirobranchus triqueter encrusts stones, rocks and shells, and the carapace of some species of decapods. They are predominantly sublittoral to depths of 70 m.

↓ Depth range

Up to 70m

Q Identifying features

- The operculum bears a shallow, dish-shaped plug (ampulla), the distal part is often conical and may have projections on the crown.
- The tube is up to 25 mm long.
- A single ridge runs along the top of the tube, ending in a projection over the anterior opening.
- Colouration of the worm is varied.

🏛 Additional information

- May be confused with *Spirobranchus lamarcki*, the tube of which differs from *Spirobranchus triqueter* as it has two vestigial ridges, one on each side, in addition to the median keel. Further differences can only be seen when the worm is removed from its tube (Hayward & Ryland, 1995). Further distinction between the two species can be obtained by using biochemical genetics, as described by Ekaratne *et al.* (1982).
- Males are cream in colour whilst females are bright pink/orange in colour (Thomas, 1940).

✓ Listed by

🔗 Further information sources

Search on:

    NBN WoRMS

Biology review

Taxonomy

Phylum	Annelida	Segmented worms e.g. ragworms, tubeworms, fanworms and spoon worms
Class	Polychaeta	Bristleworms, e.g. ragworms, scaleworms, paddleworms, fanworms, tubeworms and spoon worms
Order	Sabellida	
Family	Serpulidae	
Genus	Spirobranchus	
Authority	(Linnaeus, 1758)	
Recent Synonyms	Pomatoceros triqueter	

Biology

Typical abundance	No information found
Male size range	up to 25mm
Male size at maturity	
Female size range	Small(1-2cm)
Female size at maturity	
Growth form	Vermiform segmented
Growth rate	1.5mm/month
Body flexibility	High (greater than 45 degrees)
Mobility	
Characteristic feeding method	Active suspension feeder, No information
Diet/food source	
Typically feeds on	Plankton and detritus
Sociability	
Environmental position	Epibenthic
Dependency	Independent.
Supports	No information
Is the species harmful?	No No text entered

Biology information

Growth

- Once settled onto the substratum the worm forms a temporary delicate semi-transparent tube, which, when calcareous material is later added at the anterior end (Hayward & Ryland, 1995) dissolves over time (Dons, 1927). The tube is formed by a secretion of calcium carbonate (obtained from sea water) from the collar (Thomas, 1940).
- Growth rate is usually measured by the increase in length of the tube over a period of time. Dons (1927) found that the youngest sessile stages of the animals in Trondheim occurred when the tubes were 800-1200µm long and the animal was approximately 500µm

in length.

- Hayward & Ryland (1995) and Dons (1927) stated that growth is rapid and sexual maturity is reached in approximately 4 months. Growth rate has been observed by Dons (1927) to be 1.5 mm per month, although this varied with external conditions. Males and females exhibit the same growth rate (Castric-Fey, 1983). Animals settling during spring show the best growth rate and the rate is greatest during the first year (Castric-Fey, 1983).
- Castric-Fey (1983) reported that the number of segments of the worm increases with age, with a linear relationship being present within the first 6 months.

Feeding & Respiration

Thomas (1940) reviewed feeding and respiration in the polychaete. *Spirobranchus* (as *Pomatoceros*) *triqueter* never leaves its tube. Occasionally the posterior end of the tube becomes blocked by a calcareous plate with holes in. Respiration and excretion take place using cilia action to set up currents, bringing water in and down the length of the tube and flushing it back out the same way. Respiration occurs through the surface of the body and the branchial crown.

Feeding takes place by spreading apart its branchial filaments to expose a central groove. Using cilia action, it induces a current and transports food particles towards its mouth. If particles are too large or too numerous, the tip of a filament bends over and removes it. No sorting of food particles takes place.



Habitat preferences

Physiographic preferences	Enclosed coast / Embayment, Open coast
Biological zone preferences	Lower infralittoral, Sublittoral fringe, Upper circalittoral, Upper infralittoral
Substratum / habitat preferences	Artificial (man-made), Bedrock, Cobbles, Crevices / fissures, Gravel / shingle, Large to very large boulders, Pebbles, Small boulders
Tidal strength preferences	Moderately Strong 1 to 3 knots (0.5-1.5 m/sec.), Strong 3 to 6 knots (1.5-3 m/sec.), Very Weak (negligible), Weak < 1 knot (<0.5 m/sec.)
Wave exposure preferences	Exposed, Extremely sheltered, Moderately exposed, Sheltered, Very exposed, Very sheltered
Salinity preferences	Full (30-40 psu)
Depth range	Up to 70m
Other preferences	No text entered
Migration Pattern	Non-migratory / resident

Habitat Information

- Segrove (1941) studied *Spirobranchus triqueter* in south England and found that there are usually ten times as many males as females present.
- The species has been noted to occur in very exposed to extremely sheltered wave action, very sheltered to exposed water flow rate, and in areas where there is little or no silt present (Price *et al.*, 1980).
- *Spirobranchus triqueter* is considered to be a primary fouling organism (Crisp, 1965),

colonizing artificial commercially important structures such as buoys, ships hulls, docks and offshore oil rigs (OECD, 1967).

- *Spirobranchus triqueter* is an opportunistic species, making use of available space quickly. In Bantry Bay, south-west Ireland, fouling by the tube worm caused a 65% mortality of scallops and prevented scallops from recolonizing the area after spat collection (Burnell *et al.*, 1991). They also reported that mussel farmers considered that most inner areas of the bay would be subject to this type of fouling.
- Rubin (1985) reported that *Spirobranchus* (syn. *Pomatoceros*) *triqueter* overgrew colonies of encrusting Bryozoa to become the dominant species on experimental panels. However, Bryozoa then grew on the tubes of the species, thereby avoiding exclusion.
- Dominance of *Spirobranchus lamarckii* over *Spirobranchus triqueter* is dependent on climatic conditions (Castric-Fey, 1983).

Life history

Adult characteristics

Reproductive type	Protandrous hermaphrodite
Reproductive frequency	Annual episodic
Fecundity (number of eggs)	No information
Generation time	Insufficient information
Age at maturity	Approximately 4 months
Season	See additional information
Life span	See additional information

Larval characteristics

Larval/propagule type	-
Larval/juvenile development	Planktotrophic
Duration of larval stage	11-30 days
Larval dispersal potential	Greater than 10 km
Larval settlement period	See additional information

Life history information

- Male *Spirobranchus triqueter* release spermatogonia or primary spermatocytes and females release primary oocytes through a pair of gonoducts, consisting of a ciliated funnel and tube (Thomas, 1940).
- Hayward & Ryland (1995) and Segrove (1941) suggested that breeding of *Spirobranchus triqueter* probably takes place throughout the year. However, Hayward & Ryland (1995) noted a breeding peak in spring and summer and records from Port Erin by Moore (1937) indicated that breeding only took place in April in this location.
- Castric-Fey (1983) studied variations in settlement rate and concluded that, although the species settled all year round, very rare settlement was observed during winter and maximum settlement occurred in April, June, August and Sept-Oct. Studies in Bantry Bay (Cotter *et al.*, 2003) revealed a single peak in recruitment during summer (especially July

and August) with very little recruitment at other times of the year. More individuals settled on panels at 7 m than at 4 m.

- Larvae are pelagic for about 2-3 weeks in the summer. However, in the winter this amount of time increases to about 2 months (Hayward & Ryland, 1995).

Longevity

Longevity has been recorded to be between 1.5 to 4 years. Hayward & Ryland (1995) noted that individuals lived approximately 1.5 years, with most individuals dying after breeding (Hayward & Ryland, 1995). Castric-Fey (1983) found that under laboratory conditions, individuals were still alive after 2.5 years. However, Castric-Fey (1983) also stated that under natural conditions it is probable that they do not live any longer than this. Whilst Dons (1927) found that, according to measured growth rate, some of the individuals he studied would have been at least 4 years old.

Sensitivity review

This MarLIN sensitivity assessment has been superseded by the MarESA approach to sensitivity assessment. MarLIN assessments used an approach that has now been modified to reflect the most recent conservation imperatives and terminology and are due to be updated by 2016/17.

A Physical Pressures

	Intolerance	Recoverability	Sensitivity	Confidence
Substratum Loss	High	High	Moderate	High

Pomatoceros triqueter is attached permanently to rocks, boulders or shingle. Removal of substratum will remove calcareous tubes and animals contained in them. Intolerance is assessed as high. Recoverability is likely to be high (see additional information below).

Smothering	High	High	Moderate	High
-------------------	------	------	----------	------

Smothering with a 5 cm layer of sediment would completely cover the tubes of *Pomatoceros triqueter* that usually lie flat against the surface of the rock. It is also likely that too much sediment on the surface of rocks or shells would prevent settlement of larvae and impair the long term survival of populations. Intolerance has been assessed to be high. Recoverability is likely to be high (see additional information below).

Increase in suspended sediment	Low	High	Low	Low
---------------------------------------	-----	------	-----	-----

Records show confusion as to whether *Pomatoceros triqueter* is tolerant of high suspended sediment levels. According to Bacescu (1972), sabellids are accustomed to turbidity and silt. Stubbings & Houghton (1964) found *Pomatoceros triqueter* in Chichester harbour, a muddy harbour, therefore agreeing with the previous statement. However, *Pomatoceros triqueter* has been noted to occur in areas where there is little or no silt present (Price *et al.*, 1980) and according to Lewis (1957), *Pomatoceros triqueter* is highly susceptible to unfavourable conditions, always requiring stability and clean water. Moore (1937) and Nair (1962) agreed with this.

However, *Pomatoceros triqueter* has been recorded in areas where suspended sediment levels can be high; demonstrating that it can tolerate high suspended sediment concentrations. A supply of suspended sediment will probably also be important to *Pomatoceros triqueter* because the species requires a supply of particulate matter for suspension feeding. At the benchmark level of an increase of 100mg/l for one month, the likely impact would be an increase in cleaning costs. Intolerance has been assessed as low. Recoverability is likely to be high.

Decrease in suspended sediment	Low	High	Low	Moderate
---------------------------------------	-----	------	-----	----------

Pomatoceros triqueter has been noted to occur in areas where there is little or no silt present (Price *et al.*, 1980). The species is an active suspension feeder and will probably not be highly intolerant of suspended sediment concentrations. As an energetic cost would probably be entailed to create currents to transport food particles, intolerance has been assessed to be low. On return to normal conditions, recoverability is likely to be high.

Dessication	Intermediate	High	Low	Moderate
--------------------	--------------	------	-----	----------

As *Pomatoceros triqueter* occurs in the subtidal region it will be tolerant to a certain amount of desiccation. The species probably survives by closing the operculum of the tube, however, the

amount of time available for feeding and respiration will be reduced, and therefore the population's viability may be reduced. Some individuals are also likely to die. Intolerance has been assessed to be intermediate. Recoverability is likely to be high (see additional information below).

Increase in emergence regime Intermediate High Low Low

An increase in the emergence regime will increase the amount of time some individuals are exposed to air. At the benchmark level of an increase of one hour over the period of a year, those higher on shore are likely to die. Intolerance has been assessed to be intermediate. Recoverability is likely to be high (see additional information below).

Decrease in emergence regime Tolerant* Not relevant Not sensitive* Not relevant

A decrease in the emergence regime may mean that more time can be spent feeding, but is unlikely to have any adverse effects. Therefore *Pomatoceros triqueter* is likely to tolerate a decrease in emergence, and may actually benefit.

Increase in water flow rate Tolerant* Not relevant Not sensitive* Not relevant

Pomatoceros triqueter has been noted to occur in areas with very sheltered to exposed water flow rates (Price *et al.*, 1980). Wood (1988) observed *Pomatoceros* sp. in strong tidal streams and Hiscock (1983) found that in strong tidal streams or strong wave action where abrasion occurs, fast growing species such as *Pomatoceros triqueter* occur. Therefore, the species is probably tolerant of an increase in water flow rate, and the species may actually increase in abundance.

Decrease in water flow rate Tolerant Not relevant Not sensitive Not relevant

Pomatoceros triqueter has been noted to occur in areas with very sheltered to exposed water flow rates (Price *et al.*, 1980). The species has been assessed to be tolerant.

Increase in temperature Tolerant* Not relevant Not sensitive* Not relevant

Maximum sea surface temperatures around the British Isles rarely exceed 20 °C (Hiscock, 1998) and, as *Pomatoceros triqueter* occurs as far south as the Mediterranean, it will therefore be subject to a wider range of temperatures than experienced in the British Isles. Further information also backs this up. Castric-Fey (1983) found that animals settling during spring showed the best growth rate and the best larval settlement occurred in the summer months. *Pomatoceros triqueter* has been assessed as tolerant* to an increase in temperature.

Decrease in temperature Intermediate High Low Moderate

Minimum surface sea water temperatures rarely fall below 5 °C around the British Isles (Hiscock, 1998). Below a temperature of 7°C *Pomatoceros triqueter* is unable to build calcareous tubes (Thomas, 1940). This means that, although adults may be able to survive a decrease in temperature, larvae would not be able to attach to the substratum. Intolerance has been assessed to be intermediate. Recoverability is likely to be high (see additional information below).

Increase in turbidity Tolerant Not relevant Not sensitive Not relevant

According to Bacescu (1972), sabellids are accustomed to turbidity and silt. *Pomatoceros triqueter* has also recently been recorded by De Kluijver (1993) from Scotland in the aphotic zone, indicating that the species would not be sensitive to an increase in turbidity.

Decrease in turbidity Tolerant Not relevant Not sensitive Not relevant

According to Bacescu (1972), sabellids are accustomed to turbidity and silt. According to

Lewis (1957), *Pomatoceros triqueter* is highly susceptible to unfavourable conditions, always requiring stability and clean water. Moore (1937) and Nair (1962) agreed with this. Therefore, *Pomatoceros triqueter* is unlikely to be sensitive to a decrease in turbidity.

Increase in wave exposure

Low

High

Low

Moderate

Pomatoceros triqueter has been noted to occur in areas with variable wave action; extremely sheltered to very exposed (Price *et al.*, 1980). The hard calcareous tube is resistant to abrasion from sand, gravel and boulders (Wood, 1988; Hiscock, 1983) that are mobilised by wave action. With an increase in wave exposure over a period of a year the viability of the population may be reduced due to a reduction in feeding and larval settlement. Therefore intolerance of *Pomatoceros triqueter* to an increase in wave exposure is likely to be low. On return to normal conditions, recoverability is likely to be high.

Decrease in wave exposure

Tolerant

Not relevant

Not sensitive

Not relevant

Pomatoceros triqueter has been noted to occur in areas with variable wave action; extremely sheltered to very exposed (Price *et al.*, 1980). As the species can tolerate very low wave exposure, it is therefore probably tolerant of a decrease in wave exposure.

Noise

Tolerant

Not relevant

Not sensitive

Not relevant

Polychaetes may be able to detect vibration, and withdraw into their tube. However, at the benchmark level the species is unlikely to be sensitive to noise.

Visual Presence

Tolerant

Not relevant

Not sensitive

Not relevant

Shadows detected by the photoreceptive surface of serpulid polychaetes may result in withdrawal of the worm back into its tube (Kinne, 1970). However, at the benchmark level the species is unlikely to be sensitive to visual presence.

Abrasion & physical disturbance

Intermediate

High

Low

Not relevant

Pomatoceros triqueter has a hard calcareous tube that is resistant to sand and gravel abrasion (Wood, 1988). Hiscock (1983) noted that a community, under conditions of scour and abrasion from stones and boulders moved by storms, developed into a community consisting of fast growing species such as *Pomatoceros triqueter*. Off Chesil Bank, the epifaunal community dominated by *Pomatoceros triqueter*, *Balanus crenatus* and *Electra pilosa*, decreased in cover in October, was scoured away in winter storms, and was recolonized in May to June (Warner, 1985). Warner (1985) reported that the community did not contain any persistent individuals, being dominated by rapidly colonizing organisms. But, while larval recruitment was patchy and varied between the years studied, recruitment was sufficiently predictable to result in a dynamic stability and a similar community was present in 1979, 1980, and 1983. Scour due to winter storms is probably greater than the benchmark level. Scour and abrasion will probably remove a proportion of the population, suggesting an intolerance of intermediate. However, it demonstrates rapid growth and recruitment so that it is not considered to be sensitive. The abundance of *Pomatoceros triqueter* may increase due to decreased competition from other species.

Displacement

Low

High

Low

High

If tubes containing the worm are removed, the tubes will not be able to be reattached to the substratum surface. However, Thomas (1940) found that if *Pomatoceros triqueter* is removed from its tube, it will start to make a new one in a few hours. Therefore, it is likely that the worm will be able to leave the old tube to start constructing another. This would probably involve an added energetic cost, therefore population viability may be affected. Intolerance has been assessed to be low. Recoverability is likely to be high.

Chemical Pressures

	Intolerance	Recoverability	Sensitivity	Confidence
Synthetic compound contamination		Not relevant		Not relevant
There is insufficient information to assess the intolerance of <i>Pomatoceros triqueter</i> to synthetic chemicals.				
Heavy metal contamination		Not relevant		Not relevant
Bryan (1984) suggested that, on evidence available for several species, that polychaetes are fairly resistant to heavy metals. However, there is insufficient information available to assess intolerance of <i>Pomatoceros triqueter</i> to heavy metal contamination.				
Hydrocarbon contamination		Not relevant		Not relevant
Large numbers of dead polychaetes and other fauna were washed up at Rulosquet marsh near Isle de Grand following the Amoco Cadiz oil spill in 1978 (Cross <i>et al.</i> , 1978). However, no information was found relating to <i>Pomatoceros triqueter</i> in particular. Therefore, insufficient information was available to assess the species intolerance.				
Radionuclide contamination		Not relevant		Not relevant
There is insufficient information to assess the intolerance of <i>Pomatoceros triqueter</i> to radionuclides.				
Changes in nutrient levels		Not relevant		Not relevant
There is insufficient information to assess the intolerance of <i>Pomatoceros triqueter</i> to nutrient levels.				
Increase in salinity	Tolerant	Not relevant	Not sensitive	Not relevant
<i>Pomatoceros triqueter</i> occurs in fully saline waters and is probably relatively tolerant of an increase in salinity.				
Decrease in salinity	High	High	Moderate	Low
<i>Pomatoceros triqueter</i> occurs in fully saline coastal waters and has not been recorded from brackish or estuarine waters. Therefore, it is likely that the species will be very intolerant of a decrease in salinity. However, Dixon (1985) views the species as able to withstand significant reductions in salinity. The degree of reduction in salinity and time that the species could tolerate those levels were not recorded. Therefore, there is insufficient information available to assess the intolerance of <i>Pomatoceros triqueter</i> to a reduction in salinity.				
Changes in oxygenation		Not relevant		Not relevant
Cole <i>et al.</i> (1999) suggest possible adverse effects on marine species below 4 mg/l and probable adverse effects below 2 mg/l. However, no information was found relating to intolerance of <i>Pomatoceros triqueter</i> to oxygen levels. Insufficient information was available to assess intolerance of the species at the benchmark level of 2 mg/l for a week.				

Biological Pressures

	Intolerance	Recoverability	Sensitivity	Confidence
Introduction of microbial pathogens/parasites		Not relevant		Not relevant
Thomas (1940) recorded parasites of <i>Pomatoceros triqueter</i> . <i>Trichodina pediculus</i> (a ciliate) was observed in fair numbers moving over the branchial crown. However, this is a commensal, not				

a parasite. Parasites found in the worm include gregarines & ciliated protozoa and parasites that had the appearance of sporozoan cysts. However, no information was found about the effects of microbial pathogens on *Pomatoceros triqueter*.

Introduction of non-native species **Tolerant** Not relevant **Not sensitive** Not relevant

Although several species of serpulid polychaetes have been introduced into British waters none are reported to compete with *Pomatoceros triqueter* (Eno *et al.*, 1997).

Extraction of this species Not relevant Not relevant Not relevant Not relevant

No extraction of *Pomatoceros triqueter* is known to occur.

Extraction of other species Not relevant Not relevant Not relevant Not relevant

No extraction of other species is likely to have any effect on *Pomatoceros triqueter*.

Additional information

The species is fairly widespread, reaches sexual maturity within 4 months (Hayward & Ryland, 1995; Dons, 1927) and longevity has been recorded to be between 1.5 and 4 years (Hayward & Ryland, 1995; Castric-Fey, 1983; Dons, 1927). Larvae are pelagic for about 2-3 weeks in the summer and about 2 months in the winter (Hayward & Ryland, 1995), enabling them to disperse widely. Recovery is therefore likely to be high.

Importance review

 Policy/legislation

- no data -

 Status

National (GB) importance	-	Global red list (IUCN) category	-
--------------------------	---	---------------------------------	---

 Non-native

Native	-		
Origin	-	Date Arrived	-

 Importance information

Pomatoceros triqueter is an opportunistic species that can live on a variety of substrates; from rocks, boulders and pebbles to man-made structures. The fouling of the tube worm can compete with and exclude other species.

Bibliography

- Bacescu, M.C., 1972. Substratum: Animals. In: *Marine Ecology: A Comprehensive Treatise on Life in Oceans and Coastal Waters. Volume 1 Environmental Factors Part 3*. (ed. O. Kinne). Chichester: John Wiley & Sons.
- Bradshaw, C., Veale, L.O., Hill, A.S. & Brand, A.R., 2002. The role of scallop-dredge disturbance in long-term changes in Irish Sea benthic communities: a re-analysis of an historical dataset. *Journal of Sea Research*, **47**, 161-184.
- Bryan, G.W., 1984. Pollution due to heavy metals and their compounds. In *Marine Ecology: A Comprehensive, Integrated Treatise on Life in the Oceans and Coastal Waters*, vol. 5. *Ocean Management*, part 3, (ed. O. Kinne), pp.1289-1431. New York: John Wiley & Sons.
- Burnell, G.M., Barnett, M., O'Carroll, T. & Roantree, V., 1991. Scallop spat collection and on-growing trials in south-west Ireland. *Actes de Colloques. Instituts Francais de Recherche pour l'Exploitation de la Mer (IFREMER)*, **17**, 139-144.
- Castric-Fey, A., 1983. Recruitment, growth and longevity of *Pomatoceros triqueter* and *Pomatoceros lamarckii* (Polychaeta, Serpulidae) on experimental panels in the Concarneau area, South Brittany. *Annales de l'Institut Oceanographique, Paris*, **59**, 69-91.
- Cole, S., Codling, I.D., Parr, W. & Zabel, T., 1999. Guidelines for managing water quality impacts within UK European Marine sites. *Natura 2000 report prepared for the UK Marine SACs Project*. 441 pp., Swindon: Water Research Council on behalf of EN, SNH, CCW, JNCC, SAMS and EHS. [UK Marine SACs Project.], <http://www.ukmarinesac.org.uk/>
- Cotter, E., O'Riordan, R.M. & Myers, A.A. 2003. Recruitment patterns of serpulids (Annelida: Polychaeta) in Bantry Bay, Ireland. *Journal of the Marine Biological Association of the United Kingdom*, **83**, 41-48.
- Crisp, D.J., 1965. The ecology of marine fouling. In: *Ecology and the Industrial Society, 5th Symposium of the British Ecological Society*, 99-117 (ed. G.T. Goodman, R.W. Edwards & J.M. Lambert).
- Cross, F.A., Davis, W.P., Hoss, D.E. & Wolfe, D.A., 1978. Biological Observations, Part 5. In *The Amoco Cadiz Oil Spill - a preliminary scientific report* (ed. W.N.Ness). NOAA/EPA Special Report, US Department of Commerce and US Environmental Protection Agency, Washington.
- De Kluijver, M.J., 1993. Sublittoral hard-substratum communities off Orkney and St Abbs (Scotland). *Journal of the Marine Biological Association of the United Kingdom*, **73**, 733-754.
- Dixon, D.R., 1985. Cytogenetic procedures. *Pomatoceros triqueter*: A test system for environmental mutagenesis. In *The effects of stress and pollution in marine animals*.
- Dons, C., 1927. Om Vest og voskmåte hos *Pomatoceros triqueter*. *Nyt Magazin for Naturvidenskaberne*, **LXV**, 111-126.
- Ekaratne, K., Burfitt, A.H., Flowerdew, M.W. & Crisp, D.J., 1982. Separation of the two Atlantic species of *Pomatoceros*, *P. lamarckii* and *P. triqueter* (Annelida: Serpulidae) by means of biochemical genetics. *Marine Biology*, **71**, 257-264.
- Eno, N.C., Clark, R.A. & Sanderson, W.G. (ed.) 1997. *Non-native marine species in British waters: a review and directory*. Peterborough: Joint Nature Conservation Committee.
- Fish, J.D. & Fish, S., 1996. *A student's guide to the seashore*. Cambridge: Cambridge University Press.
- Hayward, P., Nelson-Smith, T. & Shields, C. 1996. *Collins pocket guide. Sea shore of Britain and northern Europe*. London: HarperCollins.
- Hayward, P.J. & Ryland, J.S. (ed.) 1995b. *Handbook of the marine fauna of North-West Europe*. Oxford: Oxford University Press.
- Hiscock, K., 1983. Water movement. In *Sublittoral ecology. The ecology of shallow sublittoral benthos* (ed. R. Earll & D.G. Erwin), pp. 58-96. Oxford: Clarendon Press.
- Hiscock, K., ed. 1998. *Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic*. Peterborough, Joint Nature Conservation Committee.
- Howson, C.M. & Picton, B.E., 1997. *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Belfast: Ulster Museum. [Ulster Museum publication, no. 276.]
- Kinne, O. (ed.), 1970. *Marine Ecology: A Comprehensive Treatise on Life in Oceans and Coastal Waters. Vol. 1 Environmental Factors Part 1*. Chichester: John Wiley & Sons
- Lewis, J.R., 1957. Intertidal communities of the northern and western coasts of Scotland. *Transactions of the Royal Society of Edinburgh*, **63**, 185-220.
- Moore, H.B., 1937. *Marine Fauna of the Isle of Man*. Liverpool University Press.
- Nair, N.B., 1962. Ecology of marine fouling and wood-boring organisms of western Norway. *Sarsia*, **8**, 1-88.
- OECD (ed.), 1967. *Catalogue of main marine fouling organisms*. Vol. 3: *Serpulids*. Paris: Organisation for Economic Co-operation and Development.
- Price, J.H., Irvine, D.E. & Farnham, W.F., 1980. *The shore environment. Volume 2: Ecosystems*. London Academic Press.
- Rubin, J.A., 1985. Mortality and avoidance of competitive overgrowth in encrusting Bryozoa. *Marine Ecology Progress Series*, **23**, 291-299.
- Segrove, F., 1941. The development of the serpulid *Pomatoceros triquetra* L. *Quarterly Journal of Microscopical Science*, **82**, 467-540.
- Stubbings, H.G. & Houghton, D.R., 1964. The ecology of Chichester Harbour, south England, with special reference to some

fouling species. *Internationale Revue der Gesamten Hydrobiologie*, **49**, 233-279.

Thomas, J.G., 1940. *Pomatoceros, Sabella and Amphitrite*. LMBC Memoirs on typical British marine plants and animals no.33. University Press of Liverpool. Liverpool

Wood, E. (ed.), 1988. *Sea Life of Britain and Ireland*. Marine Conservation Society. IMMEL Publishing, London

Datasets

Bristol Regional Environmental Records Centre, 2017. BRERC species records recorded over 15 years ago. Occurrence dataset: <https://doi.org/10.15468/h1ln5p> accessed via GBIF.org on 2018-09-25.

Centre for Environmental Data and Recording, 2018. IBIS Project Data. Occurrence dataset: <https://www.nmni.com/CEDaR/CEDaR-Centre-for-Environmental-Data-and-Recording.aspx> accessed via NBNAtlas.org on 2018-09-25.

Centre for Environmental Data and Recording, 2018. Ulster Museum Marine Surveys of Northern Ireland Coastal Waters. Occurrence dataset <https://www.nmni.com/CEDaR/CEDaR-Centre-for-Environmental-Data-and-Recording.aspx> accessed via NBNAtlas.org on 2018-09-25.

Cofnod – North Wales Environmental Information Service, 2018. Miscellaneous records held on the Cofnod database. Occurrence dataset: <https://doi.org/10.15468/hcgqsi> accessed via GBIF.org on 2018-09-25.

Environmental Records Information Centre North East, 2018. ERIC NE Combined dataset to 2017. Occurrence dataset: <http://www.ericnortheast.org.uk/home.html> accessed via NBNAtlas.org on 2018-09-38

Fenwick, 2018. Aphotomarine. Occurrence dataset <http://www.aphotomarine.com/index.html> Accessed via NBNAtlas.org on 2018-10-01

Fife Nature Records Centre, 2018. St Andrews BioBlitz 2014. Occurrence dataset: <https://doi.org/10.15468/erweal> accessed via GBIF.org on 2018-09-27.

Fife Nature Records Centre, 2018. St Andrews BioBlitz 2015. Occurrence dataset: <https://doi.org/10.15468/xtrbyv> accessed via GBIF.org on 2018-09-27.

Fife Nature Records Centre, 2018. St Andrews BioBlitz 2016. Occurrence dataset: <https://doi.org/10.15468/146yiz> accessed via GBIF.org on 2018-09-27.

Kent Wildlife Trust, 2018. Biological survey of the intertidal chalk reefs between Folkestone Warren and Kingsdown, Kent 2009-2011. Occurrence dataset: <https://www.kentwildlifetrust.org.uk/> accessed via NBNAtlas.org on 2018-10-01.

Kent Wildlife Trust, 2018. Kent Wildlife Trust Shoresearch Intertidal Survey 2004 onwards. Occurrence dataset: <https://www.kentwildlifetrust.org.uk/> accessed via NBNAtlas.org on 2018-10-01.

Lancashire Environment Record Network, 2018. LERN Records. Occurrence dataset: <https://doi.org/10.15468/esxc9a> accessed via GBIF.org on 2018-10-01.

Manx Biological Recording Partnership, 2018. Isle of Man historical wildlife records 1990 to 1994. Occurrence dataset: <https://doi.org/10.15468/aru16v> accessed via GBIF.org on 2018-10-01.

Merseyside BioBank., 2018. Merseyside BioBank Active Naturalists (unverified). Occurrence dataset: <https://doi.org/10.15468/smzyqf> accessed via GBIF.org on 2018-10-01.

NBN (National Biodiversity Network) Atlas. Available from: <https://www.nbnatlas.org>.

North East Scotland Biological Records Centre, 2017. NE Scotland other invertebrate records 1800-2010. Occurrence dataset: <https://doi.org/10.15468/ifjfxz> accessed via GBIF.org on 2018-10-01.

OBIS (Ocean Biogeographic Information System), 2019. Global map of species distribution using gridded data. Available from: Ocean Biogeographic Information System. www.iobis.org. Accessed: 2019-03-21

South East Wales Biodiversity Records Centre, 2018. SEWBReC Worms (South East Wales). Occurrence dataset: <https://doi.org/10.15468/5vh0w8> accessed via GBIF.org on 2018-10-02.

South East Wales Biodiversity Records Centre, 2018. Dr Mary Gillham Archive Project. Occurrence dataset: <http://www.sewbrec.org.uk/> accessed via NBNAtlas.org on 2018-10-02

Yorkshire Wildlife Trust, 2018. Yorkshire Wildlife Trust Shoresearch. Occurrence dataset: <https://doi.org/10.15468/1nw3ch> accessed via GBIF.org on 2018-10-02.